

ARTÍCULO CORTO

Primer reporte de *Anagyrus kamali* Moursi y *Gyranusoidea indica* Shafee, Alam y Agarwal (Hymenoptera: Encyrtidae), parasitoides de la cochinilla rosada del hibisco *Maconellicoccus hirsutus* (Green) (Hemiptera: Pseudococcidae) en la isla de San Andres, Colombia

First report of *Anagyrus kamali* Moursi and *Gyranusoidea indica* Shafee, Alam and Agarwal (Hymenoptera: Encyrtidae), parasitoids of the pink hibiscus mealybug *Maconellicoccus hirsutus* (Green) (Hemiptera: Pseudococcidae), on San Andres Island, Colombia

Gregory Evans¹, Takumasa Kondo², María Fernanda Maya-Álvarez³, Lilliana María Hoyos-Carvajal⁴, John Albeiro Quiroz⁴, Marcela Silva-Gómez⁴

RESUMEN

Se reporta por primera vez la presencia de *Anagyrus kamali* Moursi y *Gyranusoidea indica* Shafee, Alam y Agarwal (Hymenoptera: Encyrtidae), parasitoides de la cochinilla rosada del hibisco (CRH), *Maconellicoccus hirsutus* (Green) (Hemiptera: Pseudococcidae) en la Isla de San Andrés, Colombia. Se proveen notas breves para diferenciar las dos especies de parasitoides.

Palabras clave: enemigos naturales, control biológico, morfología, nuevos registros de distribución

ABSTRACT

Here we report for the first time the presence of *Anagyrus kamali* Moursi and *Gyranusoidea indica* Shafee, Alam and Agarwal (Hymenoptera: Encyrtidae), parasitoids of the pink hibiscus mealybug (PHM), *Maconellicoccus hirsutus* (Green) (Hemiptera: Pseudococcidae), on San Andres Island, Colombia. Brief notes are provided to allow differentiation of the two parasitoid species.

Keywords: natural enemies, biological control, morphology, new distribution record

INTRODUCTION

The pink hibiscus mealybug (PHM), *Maconellicoccus hirsutus* (Green) (Hemiptera: Pseudococcidae) (Figure 1), is now widespread throughout the Caribbean islands. The taxonomy of mealybugs is based on the morphology as seen on slide-mounted specimens studied under a compound microscope. *Maconellicoccus hirsutus* has 9-segmented antennae, cerarii usually numbering 4–6 pairs (only on posterior abdominal segments), a circulus, and oral rim tubular ducts on both dorsum and venter (Kawai, 1980; Williams, 2004; Williams and Watson, 1988). It has a huge host range and occurs in all of the zoogeographic regions of the world (Ben-Dov *et al.*, 2013). On mainland Colombia, *M. hirsutus* was originally reported from the Departments of Atlántico, Cesar, Guajira and Magdalena (Kondo *et al.*, 2008). In the Department of “San Andres, Providencia and Santa Catalina”, it was first reported in 2010 on the island of Providencia (ICA, 2010). More recently it was reported on San Andres Island where it is common on *Hibiscus* spp. and *Malvaviscus arboreus* (Kondo *et al.*, 2012).

Fecha recibido: 05/10/2012
Fecha aprobado: 05/12/2012

¹ Systematic Entomology Laboratory, United States Department of Agriculture, Animal and Plant Inspection Service, Plant Protection and Quarantine, (USDA/APHIS/PPQ), Beltsville, MD, EE.UU.

² Laboratorio de Entomología, Centro de Investigación Palmira, Corporación Colombiana de Investigación Agropecuaria (Corpoica), Palmira (Colombia).

³ Jardín Botánico de San Andrés, Universidad Nacional de Colombia, San Andrés Isla (Colombia).

⁴ Departamento Ciencias Agronómicas, Facultad de Ciencias Agrarias, Universidad Nacional de Colombia. Medellín.



Figure 1. An aggregation of *Maconellicoccus hirsutus* (Green) on leaf axils of *Hibiscus rosa-sinensis*, San Andres Island, Colombia. Photo by T. Kondo

Despite the vicinity of San Andres to Central America, the PHM was likely introduced to San Andres from mainland Colombia on ornamental plants because of the constant trade between the island and mainland Colombia (Kondo *et al.*, 2012). High populations of the PHM can result in prolific production of honeydew, subsequently causing sooty mold growth, and heavy populations may cause host-plant wilting; the mealybug injects a toxic saliva during feeding, which can cause a characteristic symptom known as bunched top in hibiscus (Hodges and Hodges, 2006).

The control of the PHM is the most recent example of a successful classical biological control program. Williams (1996) indicated that the key parasitoid of the PHM is *Anagyrus kamali* Moursi (Hymenoptera: Encyrtidae), a species that is likely of Asian origin. Reddy *et al.* (2009) reported that on the Marianna Islands, *A. kamali*, *Allotropia* sp. nr. *mecriida* (Walker) (Hymenoptera: Platygasteridae) and *Cryptolaemus montrouzieri* Mulsant (Coleoptera: Coccinellidae) have successfully suppressed populations of the PHM below the economic threshold. According to Chong (2009), the combination of natural enemies, namely, the mealybug destroyer *C. montrouzieri* and the parasitoids *A. kamali* and *Gyranusoidea indica* Shafee, Alam and Agarwal (Hymenoptera: Encyrtidae), significantly reduced the initial populations of *M. hirsutus* and maintained subsequent populations under damaging levels in the Caribbean and the USA. The parasitoid rearing program at the Division of Plant Industry in Gainesville, Florida, USA, recently ended as the parasitoids now are well established in that state (A.J. Fox, personal communication, *apud* Kondo *et al.*, 2012). *Anagyrus kamali* is such an efficient parasitoid on Puerto Rico, that mass rearing is no longer necessary because the wasp is now well established there (J.C. Rodrigues, personal communication, *apud* Kondo *et al.*, 2012). In Egypt, the encyrtid *A. kamali* is known to provide good control (Williams, 1986).

Anagyrus kamali has been recorded for various scale insects, namely: *Coccus hesperidum* L. (Coccidae), *Ferrisia virgata* (Cockerell), *Ferrisia* sp., *M. hirsutus*, *Trabutina serpentina* (Green), *Nipaeococcus viridis* (Newstead), *Nipaeococcus* sp., *Phenacoccus hirsutus* Granara de Willink, *Formicococcus robustus* (Ezzat and McConnell), *Planococcus citri* (Risso), *P. halli* Ezzat and McConnell, and *Pseudococcus* sp. (Pseudococcidae) (Noyes, 2012). However, *A. kamali* appears to be specialized on the PHM. Sagarra *et al.* (2001) conducted susceptibility, preference and suitability tests on eight mealybug species, namely, *Dysmicoccus brevipes* (Cockerell), *M. hirsutus*, *Nipaeococcus nipae* (Maskell), *P. citri*, *P. halli*, *Leptococcus neotropicus* (Williams and Granara de Willink), *Pseudococcus elisae* Borchsenius, *Puto barberi* (Cockerell), and *Saccharicoccus sacchari* (Cockerell), and determined that of the nine studied species, *M. hirsutus* was the only suitable mealybug host. According to the Chalcidoidea database (Noyes, 2012), *A. kamali* is widespread and has been reported from the Antilles, Bangladesh, Belize, Caribbean (including West Indies), Egypt, Grenada, Hawaii, India, Indonesia, Jordan, Montserrat, Pakistan, Peoples' Republic of China, Puerto Rico, Saint Lucia, Sri Lanka, St. Christopher (Kitts) and Nevis, Trinidad and Tobago, and the US Virgin Islands.

We also reared *Gyranusoidea indica* from PHM on San Andres Island. This species was described from India and has been introduced into Australia, Egypt, Guyana, Trinidad, various Caribbean countries (Noyes, 2012), and California, USA (Goolsby *et al.*, 2002). *Gyranusoidea indica* has been recorded as a parasitoid of *M. hirsutus* (Meyerdirk and Warkentin, 1999; Abd-Rabou, 2001), *F. virgata* (Cockerell) (Abd-Rabou, 2001), *N. viridis* (Newstead) (Sharaf and Meyerdirk, 1987) and *Pseudococcus longispinus* (Targioni Tozzetti) (Abd-Rabou, 2001).

MATERIALS AND METHODS

The mealybug host was identified as *M. hirsutus* by GE and TK using the taxonomic keys by Kawai (1980), Williams (2004), and Williams and Watson (1988). Scale insect hosts of *A. kamali* listed in the Chalcidoidea database (Noyes, 2012) included some older names and synonyms, thus, the currently accepted names listed in the scale insect database ScaleNet (Ben-Dov *et al.*, 2013) were used in the introduction section above.

Samples of encyrtid parasitoids were reared from the PHM feeding on twigs of *Hibiscus rosa-sinensis* L. (Malvaceae) by MFMA. The encyrtid parasitoids were identified as *A. kamali* and *G. indica* by GE using the original descriptions (Moursi, 1948; Shafee *et al.*, 1975) and additional comments by Noyes (1980) and Michaud and Evans (2000). Studied material is deposited at the United States National Museum of Natural History, Beltsville, Maryland, USA (USNM). Other material studied and collected by MSG was identified by JAQ to the genus level

using the keys by Gibson *et al.* (1997) and Noyes (1980), and is deposited at the Museo Entomológico Francisco Luis Gallego, Universidad Nacional de Colombia, Sede Medellín, Medellín, Colombia (MEFLG).

Studied material

Anagyryus kamali Moursi. Colombia: San Andres Island: 12°32'10.3"N, 81°42'30.1"W, 5 m, 08.ix.2011, coll. Maria Fernanda Maya, ex *Maconellicoccus hirsutus* on *Hibiscus rosa-sinensis* L. (Malvaceae), 1 female and 1 male on microscope slide (USNM). *Gyranusoidea indica* Shafee, Alam and Agarwal. Colombia: San Andres Island: 12°32'10.3"N, 81°42'30.1"W, 5 m, 08.ix.2011, coll. Maria Fernanda Maya, ex *M. hirsutus* on *H. rosa-sinensis* L. (Malvaceae), 1 female on microscope slide (USNM).

Other studied material

Anagyryus sp. (likely *A. kamali* Moursi). 12°32'10.3"N, 81°42'30.1"W, 5 m, 19.ix.2012, coll. Marcela Silva Gómez, ex *M. hirsutus* (on *H. rosa-sinensis* L. (Malvaceae), 60 females and 19 males in vial (MEFLG); 12°32'14.7"N, 81°42'39.6"W, 42 m, 23.ix.2012, coll. Marcela Silva Gómez, ex *M. hirsutus* on *Guazuma ulmifolia* (Malvaceae), 11 females and 13 males in vial (MEFLG); 12°34'59.1"N, 81°41'23.7"W, 2 m, 13.x.2012, coll. Marcela Silva Gómez, ex *M. hirsutus* (on *Cordia sebestena* (Boraginaceae), 1 female in vial (MEFLG).

RESULTS AND DISCUSSION

Anagyryus kamali is superficially similar to *G. indica*, but can be distinguished by having the postmarginal vein of the forewing short, less than 0.5x as long as the stigmal vein (Figure 2A); frons dark brown, flagellum yellow with first segment dark brown in the female (Figure 2B), and the male with a dark spot at the apex of the antennal club. In *G. indica*, the postmarginal vein of the forewing is long, about 2x as long as the stigmal vein (Figure 2C); frons orangish, flagellum light brown in the female (Figure 2D); the male lacks the dark spot at the apex of the antennal club. In addition, most *Anagyryus* species can be separated from *Gyranusoidea* species by having conspicuous, moderately dense white setae on the face and irregular sculpture on the frontovertex, whereas *Gyranusoidea* species have pale setae on the face, but never conspicuous, and regular, polygonal sculpture on the frontovertex. For other differences, also see Montes-Rodríguez (2012).

The two parasitoids were recently reported in two municipalities in the Department of Norte de Santander, in mainland Colombia, on *M. hirsutus* infested shoots of *H. rosa-sinensis* (Montes-Rodríguez, 2012). According to Montes-Rodríguez (2012), *A. kamali* and *G. indica* probably came to Colombia by crossing the border from Venezuela, since the two parasitoids have been produced commercially and released by individuals and personnel from state-




Figure 2. *Anagyryus kamali*: A. Wing venation. B. Antenna. *Gyranusoidea indica*: C. Wing venation. D. Antenna. Photos by G. Evans.

owned institutions in that country as a mean of biological control of the PHM (Cermeli *et al.*, 2002). In Venezuela, the two parasitoids have been released in Zulia and Táchira, states that are located along the border with Colombia (Torres *et al.*, 2007, *apud* Montes-Rodríguez, 2012).

The PHM generally becomes a pest when introduced into a new environment, and that was the case when first introduced onto the island of Providencia (ICA, 2010). However, on San Andres Island the PHM is not a serious pest, probably because the encyrtid parasitoids *A. kamali* and *G. indica* and other natural enemies are keeping it under control. The two parasitoids were probably introduced to San Andres from mainland Colombia along with *M. hirsutus*.

ACKNOWLEDGEMENTS

The authors thank the Colombian Ministry of Agriculture and Rural Development (MADR) that funded the

short-term research project: “Distribución y determinación de los niveles de daño de la cochinilla acanalada *Crypticerya multicastrices* en la isla de San Andrés y búsqueda preliminar de enemigos naturales en el Valle del Cauca”; and the Government of San Andrés, Providencia y Santa Catalina, that funded the project: “Convenio Interadministrativo de Cooperación Técnica para Montaje de Laboratorio de Investigación Científica No. 058 de 2011, entre la Gobernación Departamental y la Universidad Nacional de Colombia Sede Caribe”. Many thanks also to John Jairo Alarcon from the Instituto Colombiano Agropecuario (ICA) for kindly sending the material studied to the USNM for identification. Especial thanks to Penny Gullan (The Australian National University, Canberra, Australia), Dr. M. Bora Kaydan (Çukurova University, Adana, Turkey) and to anonymous reviewers for useful comments that helped improved the content of the manuscript.

BIBLIOGRAPHIC REFERENCES

- Abd-Rabou S. 2001. Parasitoids attack mealybugs (Homoptera: Coccoidea: Pseudococcidae) in Egypt. *Egypt J Agric Res* 79(4):1355-1376.
- Ben-Dov Y, Miller DR, Gibson GAP. 2013. ScaleNet: a database of the scale insects of the world. In: <http://www.sel.barc.usda.gov/scalenet/scalenet.htm>; accessed: October, 2012.
- Cermeli M, Morales VP, Godoy F, Romero R, Cárdenas O. 2002. Presencia de la cochinilla rosada de la cayena *Maconellicoccus hirsutus* (Green) (Hemiptera: Pseudococcidae) en Venezuela. *Entomotropica* 17(1):103-105.
- Gibson GAP, Hubner JT, Woolley JB. 1997. Annotated keys to the Genera of Nearctic Chalcidoidea (Hymenoptera). Ottawa, Canada: NCR Research Press.
- Goolsby JA, Kirk AA, Meyerdirk DE. 2002. Seasonal phenology and natural enemies of *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae) in Australia. *Fla Entomol* 85(3): 494-498.
- Hodges A, Hodges G. 2006. Pink hibiscus mealybug identification (on line). Plant Health Progress, <http://www.plantmanagementnetwork.org/sub/php/diagnosticguide/2006/hibiscus/mealybug.pdf>; accessed: October, 2012.
- ICA, Instituto Colombiano Agropecuario. 2010. Plan para el manejo y mitigación del riesgo ocasionado por la cochinilla rosada (*Maconellicoccus hirsutus*) y la chinche acanalada (*Crypticerya multicastrices*) en las islas de San Andrés y Providencia. San Andrés, Colombia: Subgerencia de protección vegetal dirección técnica de epidemiología y vigilancia fitosanitaria.
- Kawai S. 1980. Scale insects of Japan in colors. (in Japanese). Tokyo: National Agricultural Education Association.
- Kondo T, Ramos-Portilla AA, Vergara-Navarro EV. 2008. Updated list of mealybugs and putoids from Colombia (Hemiptera: Pseudococcidae and Putoidae). *Bol Mus Ent Univ Valle* 9(1):29-53.
- Kondo T, Gullan P, Ramos Portilla AA. 2012. Report of new invasive scale insects (Hemiptera: Coccoidea), *Crypticerya multicastrices* Kondo and Unruh (Monophlebidae) and *Maconellicoccus hirsutus* (Green) (Pseudococcidae), on the islands of San Andres and Providencia, Colombia, with an updated taxonomic key to iceryine scale insects of South America. *Insecta Mundi* 0265:1-17.
- Meyerdirk, DE, Warkentin RW. 1999. Non-target impact of exotic natural enemies released on *Maconellicoccus hirsutus* Green in St. Kitts, West Indies. *IOBC/WPRS Bulletin* 22(2):49-50.
- Michaud JP, Evans GA. 2000. Current status of pink hibiscus mealybug in Puerto Rico including a key to parasitoid species. *Fla Entomol* 83(1):97-101.
- Montes-Rodríguez JM. 2012. Primer registro de parasitoides de la cochinilla rosada del hibisco, *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae), en Colombia [First record of parasitoids of the pink hibiscus mealybug, *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae), in Colombia]. *Rev Colomb Entomol* 38(2):274-275.
- Moursi AA. 1948. Contributions to the knowledge of the natural enemies of mealybugs. 1. Description of two new species of *Anagyrus* (Hymenoptera : Encyrtidae). *Bull Soc Fouad Ier d'Entomol* 32:9-16.
- Noyes JS. 1980. A review of the genera of Neotropical Encyrtidae (Hymenoptera: Chalcidoidea). *Bull Brit Mus Nat Hist (Bot)* 41:197-198.
- Noyes JS. 2012. Universal Chalcidoidea Database. In: <http://www.nhm.ac.uk/chalcidoids>; accessed: October, 2012.
- Sagarra LA, Vincent C, Stewart RK. 2001. Suitability of nine mealybug species (Homoptera: Pseudococcidae) as hosts for the parasitoid *Anagyrus kamali* (Hymenoptera: Encyrtidae). *Fla Entomol* 84(1):112-116.
- Sharaf NW, Meyerdirk DE. 1987. A review on the biology, ecology and control of *Nipaecoccus viridis* (Homoptera: Pseudococcidae). *Misc Publ Entomol Soc Am* 66:1-18.
- Shafee SA, Alam M, Agarwal MM. 1975. Taxonomic survey of encyrtid parasites (Hymenoptera: Encyrtidae) in India. Aligarh Muslim University Publication, Zoological Series on Indian Insect Types 10: 22-23.
- Torres BS, Fernández I, Laya M, Soto M. 2007. Resultados del proyecto fitosanitario de prevención y control de la cochinilla rosada *Maconellicoccus hirsutus* Green, durante el período 2006 - 2007. *Entomotropica* 22(2):122.
- Williams DJ. 1986. The identity and distribution of the genus *Maconellicoccus* Ezzat (Hemiptera: Pseudococcidae) in Africa. *Bull Entomol Res* 76 (2):351-357.
- Williams DJ. 1996. A brief account of the hibiscus mealybug *Maconellicoccus hirsutus* (Hemiptera: Pseudococcidae), a pest of agriculture and horticulture, with descriptions of two related species from southern Asia. *Bull Entomol Res* 86:617-628.
- Williams DJ. 2004. Mealybugs of Southern Asia. The Natural History Museum, Kuala Lumpur: Southdene SDN, BHD.
- Williams DJ, Watson GW. 1988. The scale insects of the tropical South Pacific Region. The mealybugs (Pseudococcidae). London: CAB International Institute of Entomology.